



Contents lists available at ScienceDirect

The Journal of Arthroplasty

journal homepage: www.arthroplastyjournal.org

World Expert Meeting in Arthroplasty 2024

What Are the Contraindications for Lateral or Medial Unicompartmental Knee Arthroplasty?

Saad Tarabichi, MD ^{a,*}, Fang Rui, MD, PhD ^b, David G. Deckey, MD ^a,
 Jens T. Verhey, MD ^a, Paul Van Schuyver, MD ^a, Mohamed Rashed, MD ^c,
 Usama Saleh, MD ^d, Ali Albelooshi, MD ^e, Chuan He, MD ^f, David Jevsevar, MD ^g,
 David Musil, MD ^h, Mark J. Spanghehl, MD ^a, Joshua S. Bingham, MD ^a

^a Department of Orthopaedic Surgery, Mayo Clinic Arizona, Phoenix, Arizona^b Department of Joint Surgery, The Fourth Affiliated Hospital of Xinjiang Medical University, Urumqi, China^c Department of Orthopaedic Surgery, Ministry of Health, Cairo, Egypt^d Department of Orthopaedic Surgery, Medcare Hospital, Dubai, United Arab Emirates^e Department of Orthopaedic Surgery, Mediclinic Hospital, Dubai, United Arab Emirates^f Department of Orthopaedic Surgery, Ruijin Hospital, Shanghai, China^g Adult Reconstruction Division, OrthoVirginia, Richmond, Virginia^h Department of Orthopaedic Surgery, Hospital Ceske Budejovice, Ceske Budejovice, Czech Republic

ARTICLE INFO

Article history:

Received 19 September 2024

Received in revised form

7 October 2024

Accepted 8 October 2024

Available online xxx

Keywords:

unicompartmental knee arthroplasty

medial

lateral

contraindications

total knee arthroplasty

What are the Contraindications for Lateral or Medial Unicompartmental Knee Arthroplasty?

Recommendation: In recent years, evidence suggests that many of the originally described contraindications to unicompartmental knee arthroplasty (UKA) are no longer applicable in modern clinical practice. The current contraindications for UKA include uncontrolled inflammatory arthritis, body mass index (BMI) > 35, flexion contracture > 10 degrees, coronal deformity > 10 degrees, and lateral facet patellofemoral arthritis.

Strength of Recommendation: Moderate.

Vote Results: Agree 81.9%, Disagree 9.6%, Abstain 8.5%

Rationale

Although several surgical techniques have been described for the treatment of single-compartment knee osteoarthritis, choosing the most appropriate intervention for a young, active patient can be challenging [1]. Recent data indicate that high tibial osteotomy, once popular for this demographic, has largely been replaced by unicompartmental knee arthroplasty (UKA) in younger patients [2].

One or more of the authors of this paper have disclosed potential or pertinent conflicts of interest, which may include receipt of payment, either direct or indirect, institutional support, or association with an entity in the biomedical field which may be perceived to have potential conflict of interest with this work. For full disclosure statements refer to <https://doi.org/10.1016/j.arth.2024.10.043>.

* Address correspondence to: Saad Tarabichi, MD, Department of Orthopaedic Surgery, Mayo Clinic Arizona, 5777 E Mayo Blvd, Phoenix, AZ 85054.

Several studies show that UKA is associated with reduced morbidity, decreased blood loss, shorter hospital stays, and improved postoperative range of motion compared to primary total knee arthroplasty (TKA) [3–7]. Furthermore, recent advances in implant design and surgical technique have resulted in favorable clinical outcomes and significant improvement in implant survivorship in patients undergoing UKA [8,9]. However, not all patients who had single-compartment osteoarthritis are suitable candidates for UKA. In 1989, Kozinn and Scott were the first to describe the contraindications to receiving a UKA [10]. These include but are not limited to disease in greater than one compartment, inflammatory arthropathy, nonintact anterior cruciate ligament, lateral joint line tenderness, age less than 60 years, weight greater than 82 kilograms, preoperative range of motion less than 90 degrees, flexion

<https://doi.org/10.1016/j.arth.2024.10.043>

0883-5403/© 2024 Elsevier Inc. All rights are reserved, including those for text and data mining, AI training, and similar technologies.

contracture deformity (FCD) greater than five degrees, and coronal angular deformity greater than five degrees.

Given its association with the development of osteoporosis and osteopenia, rheumatoid arthritis was previously considered to be an absolute contraindication to UKA [11]. However, with the advent of disease-modifying antirheumatic drugs, the overall morbidity secondary to rheumatoid arthritis has been significantly reduced [12]. In a recent study, Deckey et al. found that there was no difference in 2-year revision rates in UKA patients who had rheumatoid arthritis when compared to those who did not have rheumatoid arthritis (2.6 versus 2.0%, respectively, $P = 0.310$) [13]. Similarly, it was also believed that all UKAs inevitably fail and require conversion to a TKA. Therefore, to maximize implant longevity, patients who had single-compartment disease that were younger than 60 years were typically recommended to undergo TKA. However, in a prospective study that enrolled 1,000 patients, Kennedy et al. demonstrated that, with the exception of patients above 75 years, there was no association between age and implant survivorship or functional outcomes in patients undergoing medial meniscal-bearing UKA [14]. Obesity is another comorbidity that was believed to have an impact on the success rates of UKA. Although this historically meant that UKAs were not performed in patients who had a BMI of ≥ 30 , it has now been shown that mobile and fixed-bearing UKA demonstrate excellent results in the obese patient population [15]. Notwithstanding, there are data to suggest that patients who had a BMI of ≥ 35 experience suboptimal outcomes with the use of a fixed-bearing UKA construct, emphasizing the importance of appropriate implant selection in these patients [16]. Additionally, a flexion contracture deformity of greater than five degrees was traditionally considered to be a contraindication for UKA; however, a number of studies have shown that UKA can be a viable option in patients who have FCD of up to 10 degrees. In a study by Chen et al., patients who had a preoperative FCD of greater than 10 degrees had comparable outcomes to those who had an FCD of less than 10 degrees [17]. In a different study, Purcell et al. found that even at a mean FCD of 14 degrees, there was no difference in implant survivorship between the UKA and TKA groups [18]. Furthermore, patients in the UKA group had higher overall patient-reported outcome scores at the latest follow-up when compared to those that received TKA.

Over the years, there has been substantial evidence showing that not all of the originally described contraindications to UKA are applicable in modern clinical practice. Based on recent literature, the current contraindications to UKA are uncontrolled inflammatory arthritis, BMI > 35 , flexion contracture > 10 degrees, coronal deformity > 10 degrees, and lateral facet patellofemoral arthritis.

CRedit authorship contribution statement

Saad Tarabichi: Writing – review & editing, Writing – original draft. **Fang Rui:** Conceptualization. **David G. Deckey:** Conceptualization. **Jens T. Verhey:** Conceptualization. **Paul Van Schuyver:** Conceptualization. **Mohamed Rashed:** Conceptualization. **Usama Saleh:** Conceptualization. **Ali Albelooshi:** Conceptualization.

Chuan He: Conceptualization. **David Jevsevar:** Conceptualization. **David Musil:** Conceptualization. **Mark J. Spangehl:** Conceptualization. **Joshua S. Bingham:** Conceptualization.

References

- [1] Låstad Lygre SH, Espehaug B, Havelin LI, Furnes O, Vollset SE. Pain and function in patients after primary unicompartmental and total knee arthroplasty. *J Bone Joint Surg Am* 2010;92:2890. <https://doi.org/10.2106/JBJS.I.00917>.
- [2] Song SJ, Bae DK, Kim KI, Lee CH. Conversion total knee arthroplasty after failed high tibial osteotomy. *Knee Surg Relat Res* 2016;28:89–98. <https://doi.org/10.5792/ksrr.2016.28.2.89>.
- [3] Rougraff BT, Heck DA, Gibson AE. A comparison of tricompartmental and unicompartmental arthroplasty for the treatment of gonarthrosis. *Clin Orthop Relat Res* 1991;157–64.
- [4] Chassin EP, Mikosz RP, Andriacchi TP, Rosenberg AG. Functional analysis of cemented medial unicompartmental knee arthroplasty. *J Arthroplasty* 1996;11:553–9. [https://doi.org/10.1016/s0883-5403\(96\)80109-4](https://doi.org/10.1016/s0883-5403(96)80109-4).
- [5] Laurencin CT, Zelicof SB, Scott RD, Ewald FC. Unicompartmental versus total knee arthroplasty in the same patient. A comparative study. *Clin Orthop Relat Res* 1991;151–6.
- [6] Liddle AD, Pandit H, Judge A, Murray DW. Patient-reported outcomes after total and unicompartmental knee arthroplasty: a study of 14,076 matched patients from the National Joint Registry for England and Wales. *Bone Joint J* 2015;97-B:793–801. <https://doi.org/10.1302/0301-620X.97B6.35155>.
- [7] Walker T, Streit J, Gotterbarm T, Bruckner T, Merle C, Streit MR. Sports, physical activity and patient-reported outcomes after medial unicompartmental knee arthroplasty in young patients. *J Arthroplasty* 2015;30:1911–6. <https://doi.org/10.1016/j.arth.2015.05.031>.
- [8] Carlson SW, Lu Y, Sierra RJ. Minimum 10-year survivorship of mobile-bearing unicompartmental arthroplasty: single surgeon, north American non-designer consecutive series. *J Arthroplasty* 2022;37:S88–93. <https://doi.org/10.1016/j.arth.2022.02.066>.
- [9] Callaghan JJ. Unicompartmental knee replacement: introduction: where have we been? Where are we now? Where are we going? *Clin Orthop Relat Res* 2005;430:272. <https://doi.org/10.1097/01.blo.0000151844.03672.0b>.
- [10] Kozinn SC, Scott R. Unicompartmental knee arthroplasty. *J Bone Joint Surg Am* 1989;71:145–50.
- [11] Kareem R, Botleroo RA, Bhandari R, Ogeyingbo OD, Ahmed R, Gyawali M, et al. The Impact of Rheumatoid Arthritis on Bone Loss: Links to Osteoporosis and Osteopenia. *Cureus* 2021;13:e17519. <https://doi.org/10.7759/cureus.17519>.
- [12] Rodriguez-Rodriguez L, Leon L, Ivorra-Cortes J, Gómez A, Lamas JR, Pato E, et al. Treatment in rheumatoid arthritis and mortality risk in clinical practice: the role of biologic agents. *Clin Exp Rheumatol* 2016;34:1026–32.
- [13] Deckey DG, Boddu SP, Christopher ZK, Spangehl MJ, Clarke HD, Gililand JM, et al. Rheumatoid arthritis is not a contraindication to unicompartmental knee arthroplasty. *J Arthroplasty* 2024;39:2003–2006.e1. <https://doi.org/10.1016/j.arth.2024.02.067>.
- [14] Kennedy JA, Matharu GS, Hamilton TW, Mellon SJ, Murray DW. Age and outcomes of medial meniscal-bearing unicompartmental knee arthroplasty. *J Arthroplasty* 2018;33:3153–9. <https://doi.org/10.1016/j.arth.2018.06.014>.
- [15] Woo YL, Chen YQJ, Lai MC, Tay KJD, Chia S-L, Lo NN, et al. Does obesity influence early outcome of fixed-bearing unicompartmental knee arthroplasty? *J Orthop Surg* 2017;25:2309499016684297. <https://doi.org/10.1177/2309499016684297>.
- [16] Foo WYX, Liow MHL, Chen JY, Tay DKJ, Lo NN, Yeo SJ. All-polyethylene unicompartmental knee arthroplasty is associated with increased risks of poorer knee society knee score and lower satisfaction in obese patients. *Arch Orthop Trauma Surg* 2022;142:3977–85. <https://doi.org/10.1007/s00402-021-04325-w>.
- [17] Chen JY, Loh B, Woo YL, Chia SL, Lo NN, Yeo SJ. Fixed flexion deformity after unicompartmental knee arthroplasty: how much is too much. *J Arthroplasty* 2016;31:1313–6. <https://doi.org/10.1016/j.arth.2015.12.003>.
- [18] Purcell RL, Cody JP, Ammeen DJ, Goyal N, Engh GA. Elimination of preoperative flexion contracture as a contraindication for unicompartmental knee arthroplasty. *J Am Acad Orthop Surg* 2018;26:e158–63. <https://doi.org/10.5435/JAAOS-D-16-00802>.